

Esophageal Squamous Cell Carcinoma Recurring as a Solitary Renal Mass

Do Hyoung Lim, M.D.¹, Young-Hyuck Im, M.D.¹, Sang Hoon Ji, M.D.¹, Byeong-Bae Park, M.D.¹, Mi Jung Oh, M.D.¹, Jeeyun Lee, M.D.¹, Keun Woo Park, M.D.¹, Se-Hoon Lee, M.D.¹, Joon-Oh Park, M.D.¹, Kihyun Kim, M.D.¹, Won Seog Kim, M.D.¹, Chul Won Jung, M.D.¹, Young Suk Park, M.D.¹, Won Ki Kang, M.D.¹, Mark H Lee, M.D.¹, Kwanmien Kim, M.D.², Young Mog Shim, M.D.² and Keunchil Park, M.D.¹

¹Division of Hematology-Oncology, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea, ²Division of Thoracic Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Herein, a case of solitary, unilateral renal metastasis in a patient with curatively resected thoracic esophageal carcinoma, who achieved a pathological complete remission after neoadjuvant concurrent chemoradiotherapy, is reported. The kidney is the 4th or 5th most common visceral metastasis site of a primary esophageal carcinoma. More than 50% of renal metastases typically show bilateral involvement. Solitary, unilateral renal metastasis is extremely rare. Renal metastases from a primary esophageal carcinoma are usually latent and its diagnosis is very unusual in a live patient. The solitary renal

metastasis in this case was not accompanied by metastases to other sites. The value of a nephrectomy in solitary renal metastasis of esophageal cancer is not known due to the rarity of such cases. A nephrectomy could be justified in limited situations, such as with uncertainty of histological diagnosis, severe life-threatening hematuria, which cannot be controlled by embolization, or solitary renal metastasis with a long disease-free interval. (*Cancer Research and Treatment 2004;36:271-274*)

Key Words: Esophageal carcinoma, Metastasis, Kidney

INTRODUCTION

An esophageal carcinoma is often diagnosed at an advanced stage and only a small number of patients can be considered for curative resection and thus, for long-term survival (1). Surgical resection remains the only curative treatment modality for an esophageal carcinoma without distant metastasis (2). Therefore, it is important to detect distant metastases in patients with an esophageal carcinoma, so that unwarranted surgery can be avoided. Despite the increase in radical surgery for esophageal carcinomas, about 43 ~ 63% of patients eventually experience a recurrent disease after surgery (3,4). Thirty to forty percent of recurrences are due to hematogenous spread, and common sites of recurrence are the liver, lung, bone and adrenal glands. The kidney is known to be the 4th or 5th most common visceral metastasis site of esophageal carcinomas (5 ~ 7). However, solitary renal metastasis of an esophageal squamous cell carcinoma has rarely been reported. Herein, a case

of solitary, unilateral renal metastasis in a patient with curatively resected, thoracic esophageal carcinoma, who achieved a complete remission after preoperative concurrent chemoradiotherapy, is reported.

CASE REPORT

A 64-year-old man was admitted with an evaluation of progressive dysphagia in November, 2001. He had complained of dysphagia for the previous 2 months. An endoscopy showed a circular ulcerofungating mass measuring 4 cm in length, 32 cm from the incisor. A biopsy revealed a moderately differentiated, invasive squamous cell carcinoma. Staging work-up, including endoscopic ultrasonography, chest CT, abdominal CT and PET scan, showed a right high paratracheal lymph node enlargement, without distant metastasis. Under the diagnosis of locally advanced esophageal cancer, he received pre-operative concurrent chemoradiation therapy, consisting of 2 cycles of cisplatin (75 mg/m²) on day 1 and 5-fluorouracil (1,000 mg/m²) on days 1 to 4, which was repeated every 3 weeks. Radiation therapy was delivered in a total dose of 4,500 cGy in 25 fractions. He underwent an Ivor-Lewis operation, with 2-field lymph node dissection, on January 21, 2002. A histological examination showed no residual malignant cells or metastatic foci in 17 regional dissected lymph nodes, suggesting a pathologically complete remission after the preoperative chemoradiotherapy.

Correspondence: Young-Hyuck Im, Division of Hematology-Oncology, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, 50 Ilwon-dong, Gangnam-gu, Seoul 135-710, Korea. (Tel) +82-2-3410-3445, (Fax) +82-2-3410-0041, (E-mail) imyh@smc.samsung.co.kr
Received June 21, 2004, Accepted August 2, 2004




Fig. 1. CT scan of the kidney and a whole body FDG-PET scan of the patient. (A), (B) About 4.1×3.4 cm sized heterogeneous attenuating mass was detected on the anterior lip of the left kidney. (C), (D) A mass with high FDG uptake was seen on the lateral cortex area of the left kidney. The mass corresponded to the mass detected at the CT scan of the kidney.

During the regular post-operative follow-up as an outpatient, a solitary left kidney mass was detected in December, 2003. The patient was asymptomatic, with no abdominal discomfort, hematuria or any other urinary symptoms. A kidney CT scan revealed a heterogeneously attenuating mass lesion approximately 4.1×3.4 cm in size, with persistent weak enhancement on the anterior lip, mid-portion of the left kidney (Fig. 1). There were no other visceral metastases or lymphadenopathies in the CT scan. A PET scan (Fig. 1) also showed a single, abnormal increased uptake of the left kidney. He underwent a radical nephrectomy on January 14, 2004. A histological examination of the resected kidney showed a metastatic squamous cell carcinoma, which coincided with the histological findings of the previously resected esophageal carcinoma (Fig. 2). The follow-up abdominal CT scan revealed suspicious metastatic lymphadenopathy in the para-aortic space. The patient received palliative chemotherapy, consisting of cisplatin 75 mg/m² on day 1 and continuous infusion of 1,000 mg/m² 5-fluorouracil on days 1 to 4. After 3 cycles of chemotherapy, the size of the suspicious paraaortic lymphadenopathy had decreased in extent in the follow-up CT scan.

DISCUSSION

The incidence of distant metastases from a newly diagnosed esophageal carcinoma has been reported as approximately 18% (6). Several autopsy series reported that the common sites of spread as being abdominal lymph nodes (45%), followed by the liver (35%), lung (20%), cervical & supraclavicular lymph nodes (18%), bone (9%), adrenal glands (5%), peritoneum (2%) and brain (2%). Of these, the frequencies of visceral metastatic spread were as follows: lung, 22 ~ 52%; liver, 23 ~ 47%; adrenal glands, 6 ~ 20%; bone, 4 ~ 14%; kidney, 8 ~ 13% and central nervous system, 1 ~ 5% (5,8,9).

Because the renal blood flow accounts for approximately 20% of cardiac output, the kidneys are likely to be vulnerable to hematogenous metastases. More than 50% of renal metastases typically show bilateral involvement. The clinical manifestation of a metastatic renal tumor differs from that of primary renal cell carcinoma in several aspects. The majority of primary renal cell carcinomas are, on average, larger than metastatic renal tumor, with most primary tumors measuring over 6 cm at the time of presentation, compared with less than

10. Choyke PL, White EM, Zeman RK, Jaffe MH, Clark LR. Renal metastases: clinicopathologic and radiologic correlation. *Radiology* 1987;162:359-363.
 11. Heitala SO, Wahlqvist L. Metastatic tumors to the kidney: a postmortem, radiologic and clinical investigation. *Acta Radiol Diagn* 1982;23:585-591.
 12. Wada T, Ichikawa T, Okuzono T, Okuzono Y, Naito K. Renal metastasis from squamous cell carcinoma of the hypopharynx. *Urol Int* 2002;68:132-134.
 13. Yokoi H, Nakata M, Sawai K, Yoshida T, Koshikawa M, Joyama S, Tanaka A, Goto M, Ueda S, Senzaki H, Sugawara A, Kuwahara T. Intraglomerular metastasis from pancreatic cancer. *Am J Kidney Dis* 2001;37:1299-1303.
 14. Bosch A, Frias Z, Caldwell WL, Jaeschke WH. Autopsy findings in carcinoma of the esophagus. *Acta Radiol Oncol Radiat Phys Biol* 1979;18:103-112.
-