

S2 Table. Comparison of patient characteristics between group 1 and group 2 of ALBI-trend

Variable	Group 1 (n=523)	Group 2 (n=63)	p-value
Sex			
Female	221 (42.3)	17 (27.0)	0.028
Male	302 (57.7)	46 (73.0)	
Age (yr)			
< 70	414 (79.2)	39 (61.9)	0.003
≥ 70	109 (20.8)	24 (38.1)	
BMI (kg/m ²)	23.4±2.8	23.9±3.7	0.291
CEA (ng/mL)			
< 5	318 (60.8)	37 (58.7)	0.306
≥ 5	189 (36.1)	26 (41.3)	
Unknown	16 (3.1)	0	
Tumor location			
Colon	381 (72.8)	34 (54.0)	0.003
Rectum	142 (27.2)	29 (46.0)	
Tumor size (cm)			
< 5	311 (59.5)	33 (52.4)	0.345
≥ 5	212 (40.5)	30 (47.6)	
Histologic grade			
G1 & G2	484 (92.5)	56 (88.9)	0.441
G3 & MC & SRC	39 (7.5)	7 (11.1)	
LVI			
Absent	359 (68.6)	45 (71.4)	0.725
Present	153 (29.3)	16 (25.4)	
unknown	11 (2.1)	2 (3.2)	
Stage			
I & II	188 (35.9)	28 (44.4)	0.013
III	269 (51.4)	21 (33.3)	
IV	66 (12.6)	14 (22.2)	
Complications			
No	426 (81.5)	21 (33.3)	< 0.001
Yes	97 (18.5)	42 (66.7)	
Chemotherapy			
No	87 (16.6)	25 (39.7)	< 0.001
Yes	436 (83.4)	38 (60.3)	

Values are presented as number (%) or mean±SD. ALBI, albumin-bilirubin; BMI, body mass index; CEA, carcinoembryonic antigen; LVI, lymphovascular invasion; MC, mucinous adenocarcinoma; SD, standard deviation; SRC, signet-ring cell.