	Group 1 (Events/Patients)	Group 2 (Events/Patients)		HR (95% CI)	P-value
Age					
≥55	7/83	5/56	<del>-•</del>	0.86 (0.27, 2.71)	0.793
<55	3/91	6/72	<del>◆ †</del>	0.38 (0.10, 1.52)	0.172
Disease status			:		
De novo stage IV	2/61	6/33	<b>←</b> -i	0.19 (0.04, 0.93)	0.041
Recurrent	8/113	5/95	<del>-</del>	1.23 (0.40, 3.77)	0.714
Visceral disease			!		
Yes	7/97	7/75	<del>-•i</del>	0.70 (0.24, 2.00)	0.506
No	3/77	4/53	<b>◆</b> †	0.53 (0.12, 2.36)	0.404
Bone-only disease			<u> </u>		
Yes	2/32	4/40	<del>◆;</del>	0.51 (0.09, 2.79)	0.438
No	8/142	7/88	<b>-</b> ●1	0.70 (0.26, 1.95)	0.499
Number of sites			i		
≤1	3/95	4/52	<del>●  </del>	0.39 (0.09, 1.72)	0.212
2	4/42	3/43	<del>-</del>	1.13 (0.25, 5.06)	0.873
3	2/22	2/15	<del>-◆i</del>	0.58 (0.08, 4.23)	0.594
≥4	1/15	2/18	<b>-</b>	0.86 (0.08, 9.54)	0.905
			0 1 2 3 4 5 HR (95% CI)	10	
		Group 1	Better Group 2 Better		

**S3 Fig.** Forest plot of the hazard ratio for overall survival by subgroup analysis comparing group 1 with group 2 (n=302). CI, confidence interval; HR, hazard ratio.