

S1 Table. Distribution of the study population by chemotherapeutic agents

Characteristic	All	Youngest (≤ 3 yr)	Middle (3-5 yr)	Oldest (≥ 5 yr)	p-value
Chemotherapy (yes)	195 (99.5)	63 (98.4)	44 (100)	88 (100)	0.355
No. of class	5 (0-7)	4 (0-7)	6 (2-7)	5 (2-7)	< 0.001
No. of chemotherapeutic agents	7 (0-13)	5 (0-11)	8 (2-11)	7 (2-13)	< 0.001
Enzyme	73 (37.2)	15 (76.6)	28 (63.6)	30 (34.1)	< 0.001
Epipodophyllotoxin	102 (52.0)	29 (45.3)	20 (45.5)	53 (60.2)	0.117
Heavymetal	60 (30.6)	18 (28.1)	9 (20.5)	33 (37.5)	0.117
Alkylating	157 (80.1)	43 (67.2)	35 (79.5)	79 (89.8)	0.003
Antibiotics	33 (16.8)	12 (18.8)	6 (13.6)	15 (17.0)	0.782
Antimetabolites	119 (60.7)	30 (46.9)	32 (72.7)	57 (64.8)	0.015
Plantalkaloid	161 (82.1)	51 (79.7)	40 (90.9)	70 (79.5)	0.226
Corticosteroid	109 (55.6)	27 (42.2)	34 (77.3)	48 (54.5)	0.001
Anthracycline	129 (65.8)	39 (60.9)	36 (81.8)	54 (61.4)	0.040
Anthracycline dose	175 (30-500)	150 (40-500)	150 (30-367)	227 (81-495)	0.001

Values are presented as median (range) or number (%).