

**S2 Table.** Additionally adjusted postoperative prognostic factors

	Total (n=1,697)	ALND (n=1,539)	SNB (n=158)	p-value
Harvested LN				
Mean	15.16±7.44	16.26±6.90	4.39±1.75	< 0.001
N category				
N1	1,646 (97.0)	1,523 (99.0)	123 (77.9)	< 0.001
N1mi	51 (3.0)	16 (1.0)	35 (22.1)	
Chemotherapy				
No	109 (6.4)	100 (6.5)	9 (5.7)	0.131
Taxane-based	637 (37.6)	566 (36.8)	71 (44.9)	
Others	951 (56.0)	873 (56.7)	78 (49.4)	
Radiotherapy				
No	1,531 (90.4)	1,387 (90.4)	144 (91.1)	0.751
Yes	162 (9.6)	148 (9.6)	14 (8.9)	
Hormone therapy				
No	445 (26.2)	411 (26.7)	34 (21.5)	0.158
Yes	1,252 (73.8)	1,128 (73.3)	124 (78.5)	
Target agent				
No	1,595 (94.0)	1,443 (93.8)	152 (96.2)	0.219
Yes	102 (6.0)	96 (6.2)	6 (3.8)	

Values are presented as mean±standard deviation or number (%). ALND, axillary lymph node dissection; SNB, sentinel node biopsy; LN, lymph node.